

=> d his

(FILE 'HOME' ENTERED AT 15:13:01 ON 12 JAN 2006)

FILE 'MEDLINE, CAPLUS, BIOSIS, SCISEARCH' ENTERED AT 15:13:45 ON 12 JAN  
2006

L1 24946 S STOP CODON OR STOP SIGNAL  
L2 228 S L1 AND TRANSPOSON  
L3 5 S L2 AND END SEQUENCE  
L4 125 S (MODIFY OR CHANGE OR ALTER OR MUTATE OR TRUNCATE) AND END SEQ  
L5 2 S L4 AND L1  
L6 1 DUP REMOVE L5 (1 DUPLICATE REMOVED)  
L7 9 S L4 AND TRANSPOSON  
L8 3 DUP REMOVE L7 (6 DUPLICATES REMOVED)

=>

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:sssptal636kxm

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 "Ask CAS" for self-help around the clock  
NEWS 3 DEC 05 CASREACT(R) - Over 10 million reactions available  
NEWS 4 DEC 14 2006 MeSH terms loaded in MEDLINE/LMEDLINE  
NEWS 5 DEC 14 2006 MeSH terms loaded for MEDLINE file segment of TOXCENTER  
NEWS 6 DEC 14 CA/CAPLUS to be enhanced with updated IPC codes  
NEWS 7 DEC 21 IPC search and display fields enhanced in CA/CAPLUS with the  
IPC reform  
NEWS 8 DEC 23 New IPC8 SEARCH, DISPLAY, and SELECT fields in USPATFULL/  
USPAT2

NEWS EXPRESS JANUARY 03 CURRENT VERSION FOR WINDOWS IS V8.01,  
CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),  
AND CURRENT DISCOVER FILE IS DATED 19 DECEMBER 2005.  
V8.0 USERS CAN OBTAIN THE UPGRADE TO V8.01 AT  
<http://download.cas.org/express/v8.0-Discover/>

NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS INTER General Internet Information  
NEWS LOGIN Welcome Banner and News Items  
NEWS PHONE Direct Dial and Telecommunication Network Access to STN  
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that  
specific topic.

All use of STN is subject to the provisions of the STN Customer  
agreement. Please note that this agreement limits use to scientific  
research. Use for software development or design or implementation  
of commercial gateways or other similar uses is prohibited and may  
result in loss of user privileges and other penalties.

USPATALL users: See important information in the USPATFULL file banner.

\* \* \* \* \* STN Columbus \* \* \* \* \*  
FILE 'HOME' ENTERED AT 15:13:01 ON 12 JAN 2006

=> file medline caplus biosis scisearch

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'MEDLINE' ENTERED AT 15:13:45 ON 12 JAN 2006

FILE 'CAPLUS' ENTERED AT 15:13:45 ON 12 JAN 2006  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'BIOSIS' ENTERED AT 15:13:45 ON 12 JAN 2006  
Copyright (c) 2006 The Thomson Corporation

FILE 'SCISEARCH' ENTERED AT 15:13:45 ON 12 JAN 2006

=> s stop codon or stop signal

L1 24946 STOP CODON OR STOP SIGNAL

=> s l1 and transposon

L2 228 L1 AND TRANSPOSON

=> s l2 and end sequence

2 FILES SEARCHED...

L3 5 L2 AND END SEQUENCE

=>

=> d ibib abs 1-5

L3 ANSWER 1 OF 5 MEDLINE on STN

ACCESSION NUMBER: 2003372847 MEDLINE

DOCUMENT NUMBER: PubMed ID: 12907724

TITLE: The bacterial **transposon** Tn7 causes premature polyadenylation of mRNA in eukaryotic organisms: TAGKO mutagenesis in filamentous fungi.

AUTHOR: Lo Clive; Adachi Kiichi; Shuster Jeffrey R; Hamer John E; Hamer Lisbeth

CORPORATE SOURCE: Paradigm Genetics, Inc., 108 Alexander Drive, Research Triangle Park, NC 27709, USA.. clivelo@hkucc.hku.hk

SOURCE: Nucleic acids research, (2003 Aug 15) 31 (16) 4822-7. Journal code: 0411011. ISSN: 1362-4962.

PUB. COUNTRY: England: United Kingdom

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: English

FILE SEGMENT: Priority Journals

ENTRY MONTH: 200401

ENTRY DATE: Entered STN: 20030809

Last Updated on STN: 20040130

Entered Medline: 20040129

AB TAGKO is a Tn7-based transposition system for genome wide mutagenesis in filamentous fungi. The effects of **transposon** insertion on the expression of TAGKO alleles were examined in *Magnaporthe grisea* and *Mycosphaerella graminicola*. Northern analysis showed that stable, truncated transcripts were expressed in the TAGKO mutants. Mapping of the 3'-ends of TAGKO cDNAs revealed that they all contain Tn7 **end sequences**, regardless of the **transposon** orientation. Polyadenylation signals characteristic of eukaryotic genes, preceded by **stop codons** in all frames, are located in both ends of the bacterial **transposon**. Thus, TAGKO transcripts are prematurely polyadenylated, and truncated proteins are predicted to be translated in the fungal mutants. Depending on the extent of protein truncation, TAGKO mutations in HPD4 (encoding p-hydroxyphenylpyruvate dioxygenase) resulted in tyrosine sensitivity in the two fungi. Similarly, a particular *M.grisea* CBS1 (encoding cystathionine beta-synthase) TAGKO cDNA failed to complement cysteine auxotrophy in a yeast CBS mutant. TAGKO, therefore, represents a useful tool for in vivo study of truncated gene products in filamentous fungi.

L3 ANSWER 2 OF 5 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2003:837288 CAPLUS

DOCUMENT NUMBER: 139:333968

TITLE: Producing deletion derivatives of polypeptides using modified **transposon** with **stop codons** in all three reading frames

INVENTOR(S): Savilahti, Harri; Tieaho, Ville

PATENT ASSIGNEE(S): Finnzymes Oy, Finland

SOURCE: PCT Int. Appl., 37 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003087370	A1	20031023	WO 2003-FI285	20030414
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
FI 2002000746	A	20031019	FI 2002-746	20020418
US 2005208616	A1	20050922	US 2005-511327	20050330
PRIORITY APPLN. INFO.:			FI 2002-746	A 20020418
			WO 2003-FI285	W 20030414

AB The present invention describes an in vitro transposition-based methodol. for generation of deletion derivs. of polypeptides. An artificial **transposon** containing at least partly within its **transposon** ends a modification with translation **stop codons** in three reading frames is provided. In the method, transposition complexes are assembled using the modified **transposon** and essentially random integrations into the target plasmid, containing a polypeptide coding nucleic acid of interest, are recovered as a plasmid pool. Subsequent manipulation steps including restriction enzyme digestions and ligation result in pools of mutant clones from which deletion derivs. of a polypeptide coding nucleic acid of interest and its resp. deletion polypeptides could be produced.

REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 3 OF 5 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2003:616864 CAPLUS

DOCUMENT NUMBER: 139:318350

TITLE: The bacterial **transposon** Tn7 causes premature polyadenylation of mRNA in eukaryotic organisms: TAGKO mutagenesis in filamentous fungi

AUTHOR(S): Lo, Clive; Adachi, Kiichi; Shuster, Jeffrey R.; Hamer, John E.; Hamer, Lisbeth

CORPORATE SOURCE: Paradigm Genetics, Inc., Research Triangle Park, NC, 27709, USA

SOURCE: Nucleic Acids Research (2003), 31(16), 4822-4827  
CODEN: NARHAD; ISSN: 0305-1048

PUBLISHER: Oxford University Press

DOCUMENT TYPE: Journal

LANGUAGE: English

AB TAGKO is a Tn7-based transposition system for genome wide mutagenesis in filamentous fungi. The effects of **transposon** insertion on the expression of TAGKO alleles were examined in *Magnaporthe grisea* and *Mycosphaerella graminicola*. Northern anal. showed that stable, truncated transcripts were expressed in the TAGKO mutants. Mapping of the 3'-ends of TAGKO cDNAs revealed that they all contain Tn7 **end sequences**, regardless of the **transposon** orientation. Polyadenylation signals characteristic of eukaryotic genes, preceded by **stop codons** in all frames, are located in both ends of the bacterial **transposon**. Thus, TAGKO transcripts are prematurely polyadenylated, and truncated proteins are predicted to be translated in the fungal mutants. Depending on the extent of protein truncation, TAGKO mutations in HPD4 (encoding p-hydroxyphenylpyruvate dioxygenase) resulted in tyrosine sensitivity in the two fungi. Similarly, a particular *M.grisea* CBS1 (encoding cystathionine  $\beta$ -synthase) TAGKO cDNA failed to complement cysteine auxotrophy in a yeast CBS mutant. TAGKO, therefore, represents a useful tool for in vivo study of truncated gene products in filamentous fungi.

REFERENCE COUNT: 33 THERE ARE 33 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 4 OF 5 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN  
ACCESSION NUMBER: 2003:563096 BIOSIS  
DOCUMENT NUMBER: PREV200300564228  
TITLE: The bacterial **transposon** Tn7 causes premature  
polyadenylation of mRNA in eukaryotic organisms: TAGKO  
mutagenesis in filamentous fungi.  
AUTHOR(S): Lo, Clive [Reprint Author]; Adachi, Kiichi; Shuster,  
Jeffrey R.; Hamer, John E.; Hamer, Lisbeth  
CORPORATE SOURCE: Department of Botany, The University of Hong Kong, Pokfulam  
Road, Hong Kong, China  
clivelo@hkucc.hku.hk  
SOURCE: Nucleic Acids Research, (August 15 2003) Vol. 31, No. 16,  
pp. 4822-4827. print.  
ISSN: 0305-1048 (ISSN print).  
DOCUMENT TYPE: Article  
LANGUAGE: English  
ENTRY DATE: Entered STN: 3 Dec 2003  
Last Updated on STN: 3 Dec 2003

AB TAGKO is a Tn7-based transposition system for genome wide mutagenesis in  
filamentous fungi. The effects of **transposon** insertion on the  
expression of TAGKO alleles were examined in *Magnaporthe grisea* and  
*Mycosphaerella graminicola*. Northern analysis showed that stable,  
truncated transcripts were expressed in the TAGKO mutants. Mapping of the  
3'-ends of TAGKO cDNAs revealed that they all contain Tn7 **end**  
**sequences**, regardless of the **transposon** orientation.  
Polyadenylation signals characteristic of eukaryotic genes, preceded by  
**stop codons** in all frames, are located in both ends of  
the bacterial **transposon**. Thus, TAGKO transcripts are  
prematurely polyadenylated, and truncated proteins are predicted to be  
translated in the fungal mutants. Depending on the extent of protein  
truncation, TAGKO mutations in HPD4 (encoding p-hydroxyphenylpyruvate  
dioxygenase) resulted in tyrosine sensitivity in the two fungi.  
Similarly, a particular *M. grisea* CBS1 (encoding cystathionine  
beta-synthase) TAGKO cDNA failed to complement cysteine auxotrophy in a  
yeast CBS mutant. TAGKO, therefore, represents a useful tool for in vivo  
study of truncated gene products in filamentous fungi.

L3 ANSWER 5 OF 5 SCISEARCH COPYRIGHT (c) 2006 The Thomson Corporation on  
STN  
ACCESSION NUMBER: 2003:724714 SCISEARCH  
THE GENUINE ARTICLE: 712DY  
TITLE: The bacterial **transposon** Tn7 causes premature  
polyadenylation of mRNA in eukaryotic organisms: TAGKO  
mutagenesis in filamentous fungi  
AUTHOR: Lo C (Reprint); Adachi K; Shuster J R; Hamer J E; Hamer L  
CORPORATE SOURCE: Univ Hong Kong, Dept Bot, Pokfulam Rd, Hong Kong, Hong  
Kong, Peoples R China (Reprint); Univ Hong Kong, Dept Bot,  
Hong Kong, Hong Kong, Peoples R China; Paradigm Genet Inc,  
Res Triangle Pk, NC 27709 USA  
COUNTRY OF AUTHOR: Peoples R China; USA  
SOURCE: NUCLEIC ACIDS RESEARCH, (15 AUG 2003) Vol. 31, No. 16, pp.  
4822-4827.  
ISSN: 0305-1048.  
PUBLISHER: OXFORD UNIV PRESS, GREAT CLARENDON ST, OXFORD OX2 6DP,  
ENGLAND.  
DOCUMENT TYPE: Article; Journal  
LANGUAGE: English  
REFERENCE COUNT: 33  
ENTRY DATE: Entered STN: 5 Sep 2003  
Last Updated on STN: 5 Sep 2003  
\*ABSTRACT IS AVAILABLE IN THE ALL AND IALL FORMATS\*

AB TAGKO is a Tn7-based transposition system for genome wide mutagenesis  
in filamentous fungi. The effects of **transposon** insertion on  
the expression of TAGKO alleles were examined in *Magnaporthe grisea* and  
*Mycosphaerella graminicola*. Northern analysis showed that stable,  
truncated transcripts were expressed in the TAGKO mutants. Mapping of the  
3'-ends of TAGKO cDNAs revealed that they all contain Tn7 **end**  
**sequences**, regardless of the **transposon** orientation.  
Polyadenylation signals characteristic of eukaryotic genes, preceded by

**stop codons** in all frames, are located in both ends of the bacterial **transposon**. Thus, TAGKO transcripts are prematurely polyadenylated, and truncated proteins are predicted to be translated in the fungal mutants. Depending on the extent of protein truncation, TAGKO mutations in HPD4 (encoding p-hydroxyphenylpyruvate dioxygenase) resulted in tyrosine sensitivity in the two fungi. Similarly, a particular *M.grisea* CBS1 (encoding cystathionine beta-synthase) TAGKO cDNA failed to complement cysteine auxotrophy in a yeast CBS mutant. TAGKO, therefore, represents a useful tool for in vivo study of truncated gene products in filamentous fungi.